



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which the numerous and varied animal types can be made to fall into any evolutionary plan is to consider the radially symmetrical colonial coelenterates as representing the highest degree of purely biological perfection, and all other animals (except the sponges which are their equivalent) as derived from them through the appearance of various defects which had the anomalous result of leading to increased bodily efficiency.

THE CORRELATION OF TIME UNITS AND CLIMATIC CHANGES
IN PEAT DEPOSITS OF THE UNITED STATES AND EUROPE

BY ALFRED P. DACHNOWSKI

U. S. DEPT. OF AGRICULTURE, WASHINGTON, D. C.

Read before the Academy April 25, 1922

This paper is a continuation of a previous publication⁴ which contained a discussion of the evidence in peat deposits of the United States regarding climatic changes since the last Ice age.

For purposes of comparison a brief account in tabular form is presented to show certain correlations between peat deposits in this country with those of the continent of Europe.

1. A correlation of American with European *glacial drift sheets* has been made by Leverett.⁸ By considering with Leverett the early and late Wisconsin Glaciation contemporaneous with the Würm glaciation it is possible to correlate the corresponding minor glacial stages.

The three readvances of the ice sheet that produced the morainic systems of the late Wisconsin: the Valparaiso-Kalamazoo, Lake Border, and Port Huron may be regarded as corresponding to Penck and Brückner's¹⁰ cycle of mountain glaciations of the Alps: the Bühl, Gschnitz, and Daunstadia—and to those of the Scandinavian continental ice-fields: DeGeer's⁷ Dani-, Goti-, and Fini-glacial stadia.

2. *The Stratigraphic Study of Peat Deposits* in the United States has furnished valuable data as to the botanical composition of types of peat and their economic value. It has been shown also that the material formed under wet conditions differs in botanical composition and texture from that formed during dry periods, and that the study of separate and distinct layers in deposits formed within the main morainic systems during the glacial retreat serves to throw much light on the character of the past climatic changes.

Evidence is now presented by means of the position of conspicuous layers of forest peat, fibrous sedge and reed peat, and colloidal peat, to show the equivalence of peat deposits which were formed in the respective morainal belts of the United States and Europe. The greater structural complexity, it will be noted, is confined to areas within the oldest of the morainic

systems, while within the later and more recent positions of the ice border the deposits are younger in age and consist of only a few layers of peat. The lack of structural diversity in the uppermost layers of the older deposits is related probably to the distance of those deposits from the direct climatic influence of the later glacial sub-stages.

TABLE I
CORRELATION OF GLACIAL, CLIMATIC, AND LIFE STAGES SINCE THE LAST ICE-AGE
IN THE UNITED STATES

GLACIAL SUB-STAGES	GREAT LAKES STAGES	CLIMATIC CHANGES	ESTIMATED TIME	ANIMAL REMAINS	HUMAN REMAIN	VEGETATION SUCCESSION
8. Present	Lake Michi- gan 580 ft.		A.D.			Forest Spagnum bogs and marsh
7. Recent (post-gla- cial)	Post-Nippis- sing	Rising dry	—o— B.C.			
	Nippissing 530-590 Champlain Sea, Niagara Algogum 608- 609	Temperate, humid	7000-5000 B.C.		Mound builders?	Marsh
6. Port Hu- ron	Lundy-Duluth 615 ft.	Temperate		Mastodon		
morainic system		Dry				Forest
	Whittlesey- Chicago- Saginaw	Cool, humid			Flint implements	Marsh
5 Lake Bor- der	Maumee 805 ft.	Warm, dry	11000-8000 B.C.			
morainic system						Forest
4. Valparaiso	Late Wiscon- sin drift	Warm, hu- mid	18000-16000 B.C.	Mastodon		Marsh
Kalamazoo morainic system	Loess formed	Cool, windy, dry				Steppe?
		Temporary ameliora- tion				Low shrubs?
3. Bloomington		Cold, hu- mid	25000 B.C.			
morainic system						
2. Shelby- ville mo- raines		Cold	28000 B.C.			Tundra?
1. Wisconsin driftbor- der		Arctic	35000-30000 B.C.			

To the practical worker it is obvious that a better understanding of the structural differences in peat deposits implies a more correct estimate of their economic value.

The profile sections visualize the stratigraphic relationship of American and European peatland and they suggest the value of a more detailed

TABLE II
STRATIGRAPHIC FEATURES OF PEAT DEPOSITS IN THE U. S.

[illegible]

study of peat utilization in Europe in order to determine the best practices necessary to a successful solution of the peatland problem in this country on related peat deposits.

3. The common *time* relationship is confirmed by a series of correlations. It is based on the similarity between the groups of morainal systems as the great, basic divisions of time, and on the loess deposits resting upon the

TABLE III
CORRELATION OF GLACIAL, CLIMATIC, AND LIFE STAGES SINCE THE LAST ICE-AGE IN EUROPE

GLACIAL SUB-STAGES	BALTIC LAKE STAGES	CLIMATIC CHANGES	ESTIMATED TIME	ANIMAL REMAINS	HUMAN* REMAINS	VEGETATION SUCCESSION
8. Present	Baltic o. m.	Rising, dry, cool, moist	A.D.			Forest
7. Recent (Post-glacial)			B.C. 1800 B.C.- 300 A.D.		Iron age	Younger sphagnum bogs
		Warm, dry	3000-1800 B.C.			Forest Heath (beech, fir)
6. Daun	Litorina sea 0-22 m.	Warm, moist	4000-3000 B.C. 7000-5000 B.C.		Shell heaps Neolithic	Older sphagnum
Fini-glacial		Continental			culture Lake dwellers	bogs
					Azilian	Forest (oak)
5. Gschnitz			10000-8000 B.C.		Paleolithic	
Goti-glacial	Ancylus Lake 66-175 m.				culture	Marsh
						Forest (birch, pine)
4. Bühl		Severe Continental	18000-16000 B.C.	Steppe horse	Magdalen	Marsh Steppe
Dani-glacial	Yoldia sea 300 m.	Loess formed		Reindeer	culture	Dryas Steppe
		Temp. amelioration, Warm, arid				Low shrubs (Aspen willow)
3.				Mammoth Reindeer	Solutrean	Sub-arctic
2.					culture Aurignacean	Tundra
1. Würm drift border		Arctic	30000 B.C.		culture	

*Data taken in large part from Osborn (9)

Mousterian
culture

moraines of the Würm-Bühl interstadium and the early Wisconsin drift, on the glacial lake stages of the Great Lakes and the Baltic, on the series of climatic changes, and on the remarkable association of the layers of peat material in the deposits which are referable to the same glacial sub-stage; they mark the lesser divisions of time in the different geographic regions. These events were not local phenomena but well-marked stages

which occurred nearly simultaneously over a large part of northern America and northern Europe. The combined evidence suggests that it is possible to date with fair exactness the general course of events since the last Ice-age. This chronology may furnish important data bearing on different branches of science.

(a) The estimate on the duration of Post-Wisconsin glacial time in the United States is based chiefly on the recession of the waterfalls of the Niagara. The figures for the larger subdivisions fall within the estimates made by Reeds,¹² Coleman,³ Taylor,⁸ Chamberlain and Salisbury².

TABLE IV
PROFILE SECTIONS FROM PEAT DEPOSIT OF EUROPE

	XI	XII	XIII	XIV	XV	XVI	XVII
	GERMANY	ENGLAND	AUSTRIA ALPS	SWEDEN	SWEDEN	RUSSIA	RUSSIA
8.	Sphagnum	Top removed	Forest		(Forest)	(Forest)	(Forest)
7.		Sphagnum	+ Sphagnum	Sphagnum	+ Sphagnum	+ Sphagnum	+ Sphagnum
	Heath	?	Shrub forest	Forest shrub	?	Forest	Forest
6.	Sphagnum	Sphagnum	Sphagnum	Sedge-Reed	Sphagnum	Sedge	Sphagnum
	Sedge or						Hypnum
	Forest shrub	Forest	Forest	Reed	Forest shrub	Forest	Reed
5.							Sedge
	Sedge + Reed	Sedge + Reed	Reed	Macerated	Sedge	Macerated	
					Macerated		
	Reed or forest	Reed and drifted timber	Reed	Macerated			
	Macerated	Hypnum	Hypnum				
4.							
	Macerated	Macerated					
3.							
2.							
1.							

(b) The time units of measurement of Penck,¹⁰ DeGeer,⁷ Wahnschaffe¹⁵ and Brooks¹ are more or less similar with the dates delimiting the American stages.

4. The correlation of past *climates* in the United States with those of the European last glacial period has not been worked out on a meteorological basis but there is sufficient information at hand of considerable interest for comparison with the conclusions based on European evidence.

The results of the correlation agree in showing that, although the evolution of climate has been continuous, there have been several climatic waves since the last glacial period, each of relatively long duration. The climatic pulsation, resembling a climatic belt, consisted of a cold and humid period bordering the glacial readvance, followed by a warmer, drier period

in the accompanying interstadial time, with a tendency toward optimum conditions which favored the growth of forests. It is quite probable that the last climatic wave or belt, possibly in diminishing intensity, continues to shift northward at the present time.

5. In the past sixty centuries the power of nations has been moving rather steadily northward, and the higher production of agricultural crops has occurred in general near the northern limits of distribution. It is significant that this northward tendency in agricultural and social development seems to be continuing and that it appears to be correlated with practically the same course of events which is shown in Tables I to IV.

PROFILE SECTIONS OF PEAT DEPOSITS

- I, II, VI and VII from *Botanical Gazette*,⁵ 72, 57-89, 1921. (4)
- III. Profile of Kankakee Marsh near Crumstown, Indiana.
- IV. Cross Section of Dismal Swamp, Virginia, generalized from soundings and exposures along the feeder ditch and the Dismal Swamp canal.
- V. Cross section of Torrey Island in Lake Okeechobee, Florida, also near Okeelanta, about five miles south from the lake shore.
- VIII. Sounding on peat experiment sub-station near Sanilac, Michigan.
- IX. Profile along barge canal near James street bridge at Rome, N. Y.
- X. Sounding in central location of peat deposits at Chassel-Klingville, Michigan.
- XI. After Weber. (16) XII. After Rankin. (11)
- XIII. After Shreiber. (14) XIV. After Sernander. (13)
- XV. After Erdtman. (6) XVI, XVII. After Docturowski. (5)

¹ Brooks, C. E. P. Evolution of climate in North-west Europe. *Quar. J. Roy. Meteorol. Soc.*, 47, 1921 (173-194).

² Chamberlin, T. C. & Salisbury, R. D. The pleistocene or glacial period. *Geology*, Vol. 3, 1906.

³ Coleman, A. P. Length and character of the earliest interglacial period. *Bull. Geol. Soc. America*, 26, 1915 (243-245).

⁴ Dachnowski, A. P. Peat deposits and their evidence of climatic changes. *Chicago, Bot. Gaz.*, 72, 1921 (57-89).

⁵ Docturowski, V. S. Vidy torfa. (Les espèces de tourbe) *Viestnik torfianovo diela*, 2, 1915 (273-304).

⁶ Erdtman, O. G. E. Pollenanalytische Untersuchungen von Torfmooren und Marinen Sedimenten in Südwest-Schweden. *Arkiv. Bot.*, 17, 1921, No. 10, (73-79).

⁷ Geer, G. de A geochronology of the last 12000 years. *II Internat. Geol. Kong. Stockholm*, 1910, 1912 (241-253).

⁸ Leverett, F. and Taylor, F. B. The pleistocene of Indiana and Michigan and the history of the Great Lakes. *U. S. Geol. Survey, Monographs*, 53, 1915 (529).

⁹ Osborn, H. F. *Men of the Old Stone Age*, 1916.

¹⁰ Penck A. and Brückner, E. *Die Alpen im Eiszeitalter*, Leipzig. 3 vols. 1901-1909.

¹¹ Rankin, W. M. The lowland moors (mosses) of Lansdale (North Lancashire) and their development from Fens. *Types of British Vegetation*, 1911 (256-259).

¹² Reeds, C. A. Graphic projection of Pleistocene climatic oscillations. *Bull. Geol. Soc. of Amer.*, 26, 1915 (106-109).

¹³ Sernander, R. On the evidences of Postglacial changes of climate furnished by the peat-mosses of Northern Europe. *Geolog. Föreningens förhandl., Stockholm*, 30,

1909 (465-473), also *Congr ss G ologigues International*, 11 S ssion, Stockholm, 1910, 1, 1912 (203-211).

¹⁴ *Schreiber, H.* Nacheiszeitliche Klima nderung und Moorbildung in Skandinavien. *Oesterreichische Moorzeitschrift*, 15, 1914 (104).

¹⁵ *Wahnschaffe, F.*  ber die Gliederung der Glazialbildungen Norddeutschlands und die Stellung des norddeutschen Randlosses. *Zs. Gletscherk*, 5, 1911 (321-338).

¹⁶ *Weber, C. Z.* Was lehrt der Aufbau der Moore Norddeutschlands  ber den Wechsel des Klimas in post glacialer Zeit? *Zs. Deut. Geol. Gesell.*, 62, 1910 (143-162).

THE SPECTRAL ENERGY DISTRIBUTION AND OPACITY OF WIRE EXPLOSION VAPORS

BY J. A. ANDERSON

MOUNT WILSON OBSERVATORY, CARNEGIE INSTITUTION OF WASHINGTON

Communicated, May 22, 1922

It has been shown¹ that a fine wire, through which is passed the discharge of a large high voltage condenser, vaporizes so rapidly that the phenomenon is best described as an explosion. The actual time required for the wire to pass from the solid to the vapor state is probably less than one millionth of a second. If the explosion is made to take place in the space between two parallel planes from 2 to 3 millimeters apart, such as a slot in a block of wood, its spectrum is continuous, the lines of the metal exploded appearing as absorption lines. The maximum intensity of the light is very great, being of the order of that to be expected from a black body at a temperature of about 20,000° K.

A preliminary study of the spectral energy distribution given by the explosion of iron wires has been made, using a quartz spectrograph with a vacuum thermo-couple² and galvanometer. The value of $E\lambda d\lambda$ increases very rapidly from the infra red to about $\lambda 4300$; from this point to the ultra violet the great groups of iron absorption lines cause the values of $E\lambda d\lambda$ to oscillate considerably but its general tendency is upwards and the highest values are reached between $\lambda 2600$ and $\lambda 2150$. From $\lambda 2150$ to $\lambda 1990$ the values fall off rapidly, in part, perhaps, due to the absorption of the quartz lenses and prisms, and to that of the air.

Spectrograms of the explosions of the following metals have been made using a one meter focus grating spectrograph: copper, silver, gold, magnesium, zinc, cadmium, aluminum, tin, lead, tungsten, iron and nickel. All of these give continuous spectra of the same general intensity most of them appearing rather more intense than iron on account of the numerous absorption lines of the latter. Copper, silver and gold are anomalous in that the pure wires of these metals will not explode properly when placed in a slot in a block of wood, the main discharge always seeking a path around the